

Cost of reactive nitrogen release from human activities to the environment in the US

Jana Compton, US EPA (ORD NHEERL WED) Corvallis Oregon

Dan Sobota, Oregon DEQ

Michelle McCrackin, Stockholm University

Shweta Singh, Purdue University

EPA's Sustainable & Healthy Communities Research Program




Take home messages


- **First attempt in US to quantify damages associated with reactive N release to the environment.**
- **Opportunity to partner with agricultural community to maximize the benefits of food production yet minimize the loss of N to the environment.**

Nitrogen (N) inputs to US


increased 5-fold since 1900




Where are the largest human inputs?



Dominant Human N Source



What happens to the N inputs?



Our approach

- Trace N fate through the cascade
 - Source: Fossil fuel combustion, agriculture, sewage
 - Impacts: human health/social, ecosystems, agriculture, climate
- Combine N flux data with compiled data on N costs
 - \$/kg N (Compton et al. 2011; Birch et al. 2011; van Grinsven et al. 2013)


Drinking water contamination



Health effects of smog



Damages from eutrophication




Considerations


- First attempt to quantify damages from reactive N across the US
- Damage estimates are variable for many effects
- Linear scaling of effects of a kg of N
- These represent potential damages for a particular location

Costs of nitrogen pollution


- For comparison – the low and high values are associated with the EU N Assessment (from Van Grinsven *et al.* 2013 ES&T).




Anthropogenic N leakage to the environment, circa 2000



Freshwater damage costs, circa 2000




Spatial distribution of [TN] in NARS



Median = 0.604; mean = 1.214 mg N/L

Rebecca Bellmore et al. In prep.


Anthropogenic N leakage to the environment, circa 2000





kg N ha⁻¹ yr⁻¹



Watershed units
(8 digit Hydrologic Unit Codes)



Leakage reflected in local groundwater nitrate



Real Estate Transaction database
1989-2000, B. Hoppe


Idaho's Nitrate Areas of Concern

R. L. Mahler and K. E. Keith

Damages from source

Source/Sector	Damage cost (billion USD)
Agriculture	\$157.1
Fossil fuel	\$50.2
Sewage	\$2.3
Total damages from N	\$209.6
<i>Range</i>	<i>\$81-441</i>

Damages to endpoints



Other damage estimates

- **Cost of N impacts in the EU27, 2008**
 - \$97-625 billion USD (Van Grinsven et al. 2013)
- **Gross annual damages from NO_x and NH₃, 2002**
 - \$16 billion USD (Muller and Mendelsohn 2007)
- **Increased mortality associated with NH₃-derived PM_{2.5} from food export, 2006**
 - \$36 billion USD (Paulot & Jacob 2013 ES&T)

Summary

- **Human activities have increased N fixation by 5-fold in the US. 65% of N fixation is for agriculture.**
- **71% of N leaked ends up in water resources.**
- **Nitrogen damage costs are substantial - highest costs were in freshwater and coast.**

Summary (cont'd)

- **Many missing costs in our assessment, particularly for algal blooms.**
- **Findings can illustrate the range of benefits of N reductions (i.e. drinking water, air quality, coastal zone) within a place.**
- **Starting point for research connecting nutrients and damages to ecosystem goods and services.**

For more information →
Jana Compton compton.jana@epa.gov

Also see: EPA SAB Integrated nitrogen committee report 2011
EU Nitrogen Assessment 2011
International Nitrogen Initiative website

